5.1 REFERENCES

- Barot TG, Prajapati V, Patel PK, Shah N, Patel LD, (2009), A Validated RP-HPLC Method for Simultaneous Estimation of Indapamide Impurity (Methyl Nitrosoindoline) API form, International Journal of PharmTech Research, 1(4), 1287-1296.
- British pharmacopoeia volume I and II, London: HMSO Publication; 2007 ; 83
- Chaudhari BG, Patel NM, SHAH PB, (2007) Stability Indicating RP-HPLC Method for Simultaneous Determination of Atorvastatin and Amlodipine from Their Combination Drug Products. *Chem. Pharm. Bull.* 55(2) 241—246.
- Chitlange SS, Bagri K, Sakarkar DM, (2008), Stability Indicating RP- HPLC Method for Simultaneous Estimation of Valsartan and Amlodipine in Capsule Formulation. Asian J. Research Chem. 1(1), 15-18.
- Dash P, Das SN, Mahapatra SK, Kar SK, Annapurna MM, (2009), U.V. spectrophotometric method for the simultaneous estimation of Hydrochlorothiazide and Ramipril in bulk and its pharmaceutical dosage forms, Int J Pharm Sci Tech, 3(2), 25-29.
- Davidson AG. Ultraviolet-Visible absorption spectrophotometry. In Beckett AH, Stenlake JB, editors. Practical Pharmaceutical Chemistry. 4th Ed. Part-2. New Delhi: CBS Publishers; 2002. p. 275-300.
- Dhabale PN, Burade KB, Hosmani AH, Gonjari ID, Rakesh SU, Patil PR, (2009), “Development and Statistical Validation of UV Spectrophotometric Method for Estimation of Amlodipine Besylate in Tablet Dosage Form.” Arch Pharm Sci & Res., 1, 158 - 161.
- Ewing GW. Instrumental Methods of Chemical Analysis; 5th Edn; McGraw-Hill Book Company, New York, 1985, pp 1-7.
- Fernandesa N, Nimdeoa MS, Choudharia VP, Kulkarnia RR,. Pandea VV, Nikaljeb AG, (2008), dual wavelength and simultaneous equation spectrophotometric methods for estimation of atenolol and indapamide in their combined dosage form, Int. J. Chem. Sci., 6(1), 29-35.

- Harlikar JN, Amlani AM, (2003), Simultaneous Determination of Perindopril, Indapamide, Ramipril, Trandolapril in Pharmaceutical formulations using Reverse Phase Liquid Chromatography, Research Journal of Chemistry and Environment, 7 (4).
- Himabindu V, Subba R , Radhakrishnanand R, Suryanarayana M, (2007), A stability-indicating LC method for Candesartan Cilexetil, Chromatographia; 66, 7-8.
- ICH Q2 (R1): Validation of Analytical Procedures: Text and Methodology. International Conference on Harmonization of Technical Requirements for Registration of Pharmaceuticals for Human use, ICH Harmonised Tripartite Guideline, 2005.
- Ivanović D, Medenica M, Jančić B, Knežević N, Malenović A, Milić J, (2007), Validation of an analytical procedure for simultaneous determination of Hydrochlorothiazide, Lisinopril, and their impurities, *Acta chromatographica*, 18, 143-156.
- Joel GH, Lee E L. "Goodman and Gilman's "The Pharmacological Basis of Therapeutics", 10th edition, International edition, New York: The MC Graw Hill companies Inc. 2001; 843-45, 871-72, 971, 980, 984-95.
- Kamble N, Venkatchalam A, (2004), "Determination and validation of UVspectrophotometric method for simultaneous determination of lisinopril and amlodipine from tablet." Indian Drugs, 42, 170-175.
- Kondo T, Yoshida K, Yoshimura Y, Motohashi M, Tanayama S (1996), Characterization of conjugated metabolites of a new angiotensin II receptor antagonist, candesartan cilexetil, in rats by liquid chromatography/ electrospray tandem mass spectrometry following chemical derivatization, J Mas. Spectrom, 31, 873-878.
- Legorburu MJ, Alonso RM, Jiménez RM, Ortiz E, (1999), Quantitative Determination of Indapamide in Pharmaceuticals and Urine by High-Performance Liquid Chromatography with Amperometric Detection, Journal of Chromatographic Science, 37, 283-7.
- Luo J , Xiu-H W, Yu-kai H, Xinhua Z, (2004), HPLC determination of candesartan cilexetil and its related substances, Central south pharma, 02.

- Mehulkumar P, Ramesh V, Vinay kumar V, Srinivas R, Diwan PV, (2009), " Simultaneous Spectroscopic Estimation of Amlodipine Besylate and Olmesartan Medoximil in Tablet Dosage Form." Asian J. Research Chem., 2(2), 127-130.
- Parfit K. "Martindale The complete drug reference", 35th edition, Vol-I, Pharmaceutical press. 2007; 1035-36.
- Patil PR, Sachin UR, Dhabale PN, Burade KB, (2009), Simultaneous Estimation of Ramipril and Amlodipine by UV Spectrophotometric Method, Research J. Pharm. and Tech, .2 (2), 304-307.
- Patil UP, Gandhi SV, SENGAR MR, Rajmane VS, (2009), Simultaneous determination of atorvastatin calcium and telmisartan in tablet dosage form by spectrophotometry, International Journal of ChemTech Research, 1(4), 970-973.
- Ravi S Sankar, M. J Nanjan, M Vasudevan, Nahed Shaat, B Suresh, (1997), Simultaneous Estimation Of Atenolol And Amlodipine In Formulations By Reversed Phase – HPLC, Indian Journal of Pharmaceutical Sciences, 59(4), 171-173.
- Satoskar R S, Bhandarkar S D, Ainapure S S. "Pharmacology and Pharmacotherapeutics", 16th edition, Mumbai: Popular prakashan. 2003; 401-06.
- Shah NJ, Suhagia BN, Shah RR, Patel NM, (2009), HPTLC method for the simultaneous estimation of valsartan and hydrochlorothiazide in tablet dosage form, Indian J Pharm Sci, 71, 72-4.
- Singhvi I, Chaturvedi SC (1998), "Visible spectrophotometric methods for estimation of amlodipine besylate form tablets." Indian journal of pharmaceutical science, 60(5), 309-310.
- Stefan RI, Staden J., Mulaudzi L, Hassan Y, (2002), On-line simultaneous determination of S- and R-perindopril using amperometric biosensors as detectors in flow systems, Analytica Chimica Acta, 467, 189–195.
- Tarte PS, Wate SP, Khedikar PS, Satpure BT, Pawnikar G, (2008), Simultaneous Estimation of Nebivolol and Hydrochlorothiazide in Combined Tablet Dosage Form by Multicomponent Mode of Analysis, Asian J. Research Chem., 1(2), 77.

- United state pharmacopoeia-30. National formulary-25, Asian edition. 2000; 2482-83.
- Xiaoshan Z (2002), Determination of the content of candesartan cilexetil by HPLC, J china Pharm, 11.